arXiv:astro-ph/0610339v1 11 Oct 2006

Tachyon warm inflationary universe models

Ramón Herrera,[*] Sergio del Campo,[†] and Cuauhtemoc Campuzano[‡]
Instituto de Física, Pontificia Universidad Católica
de Valparaíso, Casilla 4059, Valparaíso, Chile.

(Dated: February 5, 2008)

Abstract

Warm inflationary universe models in a tachyon field theory are studied. General conditions required for these models to be realizable are derived and discussed. We describe scalar perturbations (in the longitudinal gauge) and tensor perturbations for these scenarios. We develop our models for a constant dissipation parameter Γ in one case and one dependent on ϕ in the other case. We have been successful in describing such of inflationary universe models. We use recent astronomical observations for constraining the parameters appearing in our model. Also, our results are compared with their analogous found in the cool inflationary case.

PACS numbers: 98.80.Cq

---

[*] Electronic address: ramon.herrera.a@mail.ucv.cl
[†] Electronic address: sdelcamp@ucv.cl
[‡] Electronic address: ccampuz@mail.ucv.cl

I. INTRODUCTION

It is well known that many long-standing problems of the Big Bang model (horizon, flatness, monopoles, etc.) may find a natural solution in the framework of the inflationary universe model [1, 2]. One of the success of the inflationary universe model is that it provides a causal interpretation of the origin of the observed anisotropy of the cosmic microwave background (CMB) radiation, and also the distribution of large scale structures [3, 4]. In standard inflationary universe models, the acceleration of the universe is driven by a scalar field (inflaton) with a specific scalar potential, and the quantum fluctuations associated with this field generate the density perturbations seeding the structure formation at late time in the evolution of the universe. The standard inflationary model is divided into two regimes; the slow roll and reheating epochs. In the slow roll period the universe inflates and all interactions between the inflation scalar field and any other field are typically neglected. Subsequently, a reheating period is invoked to end the period of inflation. After reheating, the universe is filled with radiation [5], and thus the universe is connected with the radiation Big-Bang phase.

Warm inflation is an alternative mechanisms for having successful inflation and avoiding the reheating period [6]. In warm inflation, dissipative effects are important during inflation, so that radiation production occurs concurrently with the inflationary expansion. The dissipating effect arises from a friction term which describes the processes of the scalar field dissipating into a thermal bath via its interaction with other fields. Also, warm inflation shows how thermal fluctuations during inflation may play a dominant role in producing the initial perturbations. In such of models, the density fluctuations arise from thermal rather than quantum fluctuations [7]. These fluctuations have their origin in the hot radiation and influence the inflaton through a friction term in the equation of motion of the inflaton scalar field [8]. Among the most attractive features of these models, warm inflation ends when the universe heats up to become radiation domination; at this epoch the universe stops inflating and "smoothly" enters a radiation dominated Big-Bang phase[6]. The matter components of the universe are created by the decay of either the remaining inflationary field or the dominant radiation field [9].

On the other hand, implications of string/M-theory to Friedmann-Robertson-Walker cosmological models have recently attracted great attention, in particular, those related to brane-antibrane configurations such as space-like branes[10]. The tachyon field associated with unstable D-branes might be responsible for cosmological inflation in the early evolution of the universe, due to tachyon condensation near the top of the effective scalar potential [11] which could also add some new form of cosmological dark matter at late times [12]. In fact, historically, as was empathized by Gibbons[13], if the tachyon condensate starts to roll down the potential with small initial velocity, then a universe dominated by this new form of matter will smoothly evolve from a phase of accelerated expansion (inflation) to an era dominated by a non-relativistic fluid, which could contribute to the dark matter specified above.

Usually, in any models of warm inflation, the scalar field which drives inflation is the standard inflaton field. As far as we know, a model in which warm inflation is driven by a tachyonic scalar field has not been yet studied. The main goal of the present work is to investigate the possible realization of a tachyonic warm inflationary universe model. In this way, we study cosmological perturbations, which are expressed in terms of different parameters appearing in our model. These parameters are constrained from the WMAP three-year data [4].

In section II, the dynamics of the tachyon warm inflationary models is obtained. In section III, the cosmological perturbations are investigated. In section IV, we use an exponential scalar potential in the high dissipation regime. There, we distinguish two cases: in the first scenario, we consider a constant dissipation rate Γ, and, in the second one, the dissipation rate is considered to be a function of the tachyonic field, ϕ. Here, we study possibles solutions of the approximated field equations. In section V we develop, in the same approximation, i.e. in the high dissipation regime, models which are characterized by a power law scale factor. Finally, in section VI, we give some conclusions.

II. WARM INFLATIONARY MODEL

As was noted by Gibbons [13], a rolling tachyon condensates in a spatially flat Friedmann-Robertson-Walker (FRW) cosmological model is described in terms of an effective fluid with energy-momentum tensor $T^{\mu}_{\nu} = diag(-\rho_{\phi}, p_{\phi}, p_{\phi}, p_{\phi})$, where the energy density, ρ_{ϕ}, and pressure, p_ϕ, associated with the tachyon field are defined by

$$\rho_\phi = \frac{V(\phi)}{\sqrt{1-\dot\phi^2}}, \tag{1}$$

and

$$p_\phi = -V(\phi)\sqrt{1-\dot\phi^2}\;, \tag{2}$$

respectively. Here, ϕ denotes the tachyon field (with unit $1/m_p$, where m_p represents the Planck mass) and $V(\phi) = V$ is the effective potential associated with this tachyon field.

The dynamics of the FRW cosmological model in the tachyonic case, in the warm inflationary scenario is described by the equations

$$H^2\; = \kappa\,[\rho_\phi + \rho_\gamma] = \kappa\left[\frac{V}{\sqrt{1-\dot\phi^2}} + \rho_\gamma\right], \tag{3}$$

$$\dot{\rho_\phi} + 3H(\rho_\phi + p_\phi) = -\Gamma\dot\phi^2 \Longrightarrow \frac{\ddot\phi}{(1-\dot\phi^2)} + 3H\;\dot\phi + \frac{V_{,\phi}}{V} = -\frac{\Gamma}{V}\sqrt{1-\dot\phi^2}\;\;\dot\phi, \tag{4}$$

and

$$\dot\rho_\gamma + 4H\rho_\gamma = \Gamma\dot\phi^2, \tag{5}$$

where $H = \dot a/a$ is the Hubble factor, a is a scale factor, ρ_γ is the energy density of the radiation field and Γ is the dissipation coefficient, with unit m_p^5. The dissipative coefficient is responsible for the decay of the tachyon scalar field into radiation during the inflationary regimen. In general, Γ can be assumed as a function of ϕ, and thus $\Gamma = f(\phi) > 0$ by the Second Law of Thermodynamics. Dots mean derivatives with respect to time, $V_{,\phi} = \partial V(\phi)/\partial\phi$, $\kappa = 8\pi/(3m_p^2)$ and we use units in which $c = \hbar = 1$.

During the inflationary era the energy density associated with the tachyonic field is the order of the potential, i.e. $\rho_\phi \sim V$, and dominates over the energy density associated with the radiation field, i.e. $\rho_\phi > \rho_\gamma$. Assuming the set of slow-roll conditions, $\dot\phi^2 \ll 1$, and $\ddot\phi \ll (3H + \Gamma/V)\dot\phi$ [6], the Friedmann equation is reduced to

$$H^2\; = \;\kappa V, \tag{6}$$

and Eq. (4) becomes

$$3H\,[\,1 + r\,]\,\dot\phi = -\frac{V_{,\phi}}{V}\; = -(\ln(V))_{,\phi}, \tag{7}$$

where r is the rate defined as

$$r = \frac{\Gamma}{3HV}, \tag{8}$$

and parameterizes the dissipation of our model. For the high (or weak) dissipation regimen, $r \gg 1$ (or $r < 1$), i.e. the dissipation coefficient Γ is much greater (or less) than the product $3\,H\,V$.

If we also considered that during tachyon warm inflation radiation production is quasi-stable, then $\dot{\rho}_\gamma \ll 4H\rho_\gamma$ and $\dot{\rho}_\gamma \ll \Gamma\dot{\phi}^2$. From Eq.(5) we obtain that the energy density of the radiation field becomes

$$\rho_\gamma = \frac{\Gamma\dot{\phi}^2}{4H} = \sigma T_r^4, \tag{9}$$

where σ is the Stefan-Boltzmann constant and T_r is the temperature of the thermal bath. Using Eqs.(7), (8) and (9) we get

$$\rho_\gamma = \sigma\, T_r^4 = \frac{r}{12\kappa}\frac{1}{(1+r)^2}\left[\frac{V_{,\phi}}{V}\right]^2. \tag{10}$$

Introducing the dimensionless slow-roll parameter

$$\varepsilon \equiv -\frac{\dot{H}}{H^2} = \frac{1}{6\kappa}\frac{1}{(1+r)}\left[\frac{V_{,\phi}}{V}\right]^2\frac{1}{V}\,, \tag{11}$$

it is possible to find a relation between the energy densities ρ_γ and ρ_ϕ given by

$$\rho_\gamma = \frac{r}{2(1+r)}\,\varepsilon\,\rho_\phi. \tag{12}$$

Here, the energy density of the tachyonic field during inflation corresponds to the potential energy, i.e. $\rho_\phi \sim V$.

The second slow-roll parameter η becomes

$$\eta \equiv -\frac{\ddot{H}}{H\dot{H}} \simeq \frac{1}{3\kappa(1+r)V}\left[\frac{V_{,\phi\phi}}{V} - \frac{1}{2}\left(\frac{V_{,\phi}}{V}\right)^2\right]. \tag{13}$$

The condition under which the tachyon warm inflation epoch could take place can be summarized with the parameter ε satisfying the relation $\varepsilon < 1$. This condition is analogous to the requirement in which $\ddot{a} > 0$. The condition given above is rewritten in terms of the densities by using Eq.(12) as

$$\rho_\phi > 2\frac{(1+r)}{r}\rho_\gamma, \tag{14}$$

which describes the epoch where tachyon warm inflation occur.

On the other hand, inflation ends when the universe heats up, at a time when $\varepsilon \simeq 1$, which implies

$$\rho_\phi \simeq 2\frac{(1+r)}{r}\rho_\gamma. \tag{15}$$

The number of e-folds at the end of inflation is given by

$$N(\phi) = -3\kappa \int_{\phi}^{\phi_f} \frac{V^2}{V_{,\phi}}(1+r)d\phi'. \tag{16}$$

In the following, the subscript i and f are used to denote the beginning and the end of inflation.

III. THE PERTURBATIONS

In this section we will describe scalar perturbations in the longitudinal gauge, and then, we will continue describing tensor perturbations.

III-1. Scalar Perturbations

By using the longitudinal gauge in the perturbed FRW metric, we write

$$ds^2 = (1+2\Phi)dt^2 - a(t)^2(1-2\Psi)\delta_{ij}dx^i dx^j, \tag{17}$$

where $\Phi = \Phi(t, \mathbf{x})$ and $\Psi = \Psi(t, \mathbf{x})$ are gauge-invariant variables introduced by Bardeen [14]. In momentum space, for the Fourier components $e^{i\mathbf{kx}}$, with k being the wave number, the following set of equations (which are derived from the perturbed Einstein field equations; we omit the subscript k here) arises:

$$\Phi = \Psi, \tag{18}$$

$$\dot{\Phi} + H\Phi = \frac{4\pi}{m_p^2}\left[-\frac{4\rho_\gamma\; a\; v}{3k} + \frac{V\dot{\phi}}{\sqrt{(1-\dot{\phi}^2)}}\,\delta\phi\right], \tag{19}$$

$$\frac{\ddot{(\delta\phi)}}{1-\dot{\phi}^2} + \left[3H + \frac{\Gamma}{V}\right]\dot{(\delta\phi)} + \left[\frac{k^2}{a^2} + (\ln(V))_{,\phi\phi} + \dot{\phi}\left(\frac{\Gamma}{V}\right)_{,\phi}\right]\delta\phi$$

$$= \left[\frac{1}{1-\dot{\phi}^2} + 3\right]\dot{\phi}\,\dot{\Phi} + \left[\dot{\phi}\frac{\Gamma}{V} - 2(\ln(V))_{,\phi}\right]\Phi, \tag{20}$$

$$(\dot{\delta\rho_\gamma}) + 4H\delta\rho_\gamma + \frac{4}{3}ka\rho_\gamma v - 4\rho_\gamma\dot{\Phi} - \dot{\phi}^2\Gamma_{,\phi}\delta\phi - \Gamma\dot{\phi}[2(\dot{\delta\phi}) - 3\dot{\phi}\Phi] = 0, \tag{21}$$

and

$$\dot{v} + 4Hv + \frac{k}{a}\left[\Phi + \frac{\delta\rho_\gamma}{4\rho_\gamma} + \frac{3\Gamma\dot{\phi}}{4\rho_\gamma}\delta\phi\right] = 0, \tag{22}$$

where v appears from the decomposition of the velocity field $\delta u_j = -\frac{iak_j}{k}\ v\ e^{i\mathbf{kx}}$ $(j = 1, 2, 3)$ [14].

Note that in the case of the scalar perturbations the tachyon and the radiation fields are interacting. Therefore, isocurvature (or entropy) perturbations are generated, besides of the adiabatic ones. This occur because warm inflation can be considered as an inflationary model with two basics fields [15, 16]. In this context, dissipative effects themselves can produce a variety of spectral ranging between red and blue [7, 15], and thus producing the running blue to red spectral suggested by WMAP three-year data[4]. We will return to this point soon.

Since what we need are the non-decreasing adiabatic and isocurvature modes on large scale $k \ll aH$, (which turn out to be a weak time dependent quantities), without loss of generality we may consistently neglect $\dot{\Phi}$ and those terms containing two-times derivatives and, combining with the slow-roll conditions, the equations for Φ, $\delta\phi$, $\delta\rho_\gamma$, and v are reduced to

$$\Phi = \frac{4\pi}{m_p^2}\left(\frac{V\dot{\phi}}{H}\right)\ \left[1 + \frac{\Gamma}{4HV} + \frac{\Gamma_{,\phi}\dot{\phi}}{48H^2V}\right]\ \delta\phi, \tag{23}$$

$$\left[3H + \frac{\Gamma}{V}\right](\dot{\delta\phi}) + \left[(\ln(V))_{,\phi\phi} + \dot{\phi}\left(\frac{\Gamma}{V}\right)_{,\phi}\right]\ \delta\phi\ = \left[\dot{\phi}\frac{\Gamma}{V} - 2(\ln(V))_{,\phi}\right]\ \Phi, \tag{24}$$

$$\delta\rho_\gamma \simeq \frac{\dot{\phi}^2}{4H}[\Gamma_{,\phi}\delta\phi - 3\Gamma\Phi] \Longrightarrow\ \frac{\delta\rho_\gamma}{\rho_\gamma} \simeq \frac{\Gamma_{,\phi}}{\Gamma}\ \delta\phi - 3\Phi, \tag{25}$$

and

$$v \simeq -\frac{k}{4aH}\left[\Phi + \frac{\delta\rho_\gamma}{4\rho_\gamma} + \frac{3\Gamma\dot{\phi}}{4\rho_\gamma}\ \delta\phi\right], \tag{26}$$

respectively.

The above equations can be solved taking ϕ as the independent variable instead of t. With the help of Eq. (7) we find

$$\left(3H + \frac{\Gamma}{V}\right)\frac{d}{dt} = \left(3H + \frac{\Gamma}{V}\right)\ \dot{\phi}\frac{d}{d\phi} = -(\ln(V))_{,\phi}\frac{d}{d\phi}, \tag{27}$$

and introducing an auxiliary function φ given by

$$\varphi = \frac{\delta\phi}{(\ln(V))_{,\phi}} \exp\left[\int \frac{1}{(3H+\Gamma/V)} \left(\frac{\Gamma}{V}\right)_{,\phi} d\phi\right], \tag{28}$$

we obtain the following equation for φ

$$\frac{\varphi_{,\phi}}{\varphi} = -\frac{9}{8}\frac{(\Gamma/V+2H)}{(\Gamma/V+3H)^2}\left[\Gamma + 4HV - \frac{\Gamma_{,\phi}(\ln(V))_{,\phi}}{12H(3H+\Gamma/V)}\right]\frac{(\ln(V))_{,\phi}}{V}. \tag{29}$$

Solving Eq.(29) and using Eq.(28) we find that

$$\delta\phi = C\,(\ln(V))_{,\phi}\exp[\Im(\phi)], \tag{30}$$

where $\Im(\phi)$ is given by

$$\Im(\phi) = -\int\left[\frac{1}{(3H+\Gamma/V)}\left(\frac{\Gamma}{V}\right)_{,\phi} + \frac{9}{8}\frac{(\Gamma/V+2H)}{(\Gamma/V+3H)^2}\left(\Gamma \cdot \right.\right.$$

$$\left.\left. + 4HV - \frac{\Gamma_{,\phi}(\ln(V))_{,\phi}}{12H(3H+\Gamma/V)}\right)\frac{(\ln(V))_{,\phi}}{V}\right]d\,\phi, \tag{31}$$

and C is a constant of integration.

In this way, the expression for the density perturbations becomes (see [17])

$$\delta_H = \frac{2}{5}\frac{m_p^2\,\exp[-\Im(\phi)]}{(\ln(V))_{,\phi}}\,\delta\phi. \tag{32}$$

We note here that in the absence of the dissipation coefficient, $\Gamma = 0$, we see that Eq.(32) is reduced to $\delta_H \sim V\delta\phi/(H\dot{\phi}) \sim H\delta\phi/\dot{\phi}$, which coincides with the expression obtained for cool inflation.

In the case of high dissipation, the dissipation parameter Γ is much greater that the product between the rate expansion H and the scalar potential, i.e. $r = \Gamma/3HV \gg 1$ and Eq.(32) becomes

$$\delta_H^2 = \frac{4}{255}\frac{m_p^4\,\exp[-2\,\widetilde{\Im}(\phi)]}{H^2\,r^2}\,\delta\phi^2, \tag{33}$$

where now $\widetilde{\Im}(\phi) := \Im(\phi)\mid_{r\gg 1}$ becomes

$$\widetilde{\Im}(\phi) = -\int\left[\frac{1}{3Hr}\left(\frac{\Gamma}{V}\right)_{,\phi} + \frac{9}{8}\left[1 - \frac{(\ln\Gamma)_{,\phi}(\ln(V))_{,\phi}}{36r\,H^2}\right](\ln(V))_{,\phi}\right]d\,\phi. \tag{34}$$

In first approximation, and during the slow-roll phase, the relation between the density matter fluctuation, $\delta\rho$, and the metric perturbation, Φ, is given by

$$\delta\rho \simeq V_{,\phi}\,\delta\phi \simeq\; -2\,[1+r]\,V\,\left[1 + \frac{\Gamma}{4HV} + \frac{\Gamma_{,\phi}\dot{\phi}}{48H^2V}\right]^{-1}\Phi, \tag{35}$$

where we have used Eq. (23). Note that in the absence of dissipation, i.e. $\Gamma = 0$, we recover the usual relation $\delta\rho/\rho \simeq -2\Phi$ valid for cool inflation, in which $\rho \simeq V$.

The fluctuations of the tachyon field are generated by thermal interaction with the radiation field, instead of quantum fluctuations. Following Ref.[18], we may write in the case $r \gg 1$,

$$(\delta\phi)^2 \simeq \frac{k_F\, T_r}{2\, m_p^4\, \pi^2}, \tag{36}$$

where the wave number k_F is defined by $k_F = \sqrt{\Gamma H/V} = H\sqrt{3r} \geq H$, and corresponds to the freeze-out scale at which dissipation damps out the thermally excited fluctuations. The freeze-out wave number k_F is defined at the point where the inequality $(\ln(V))_{,\phi\,\phi} < \Gamma H/V$, is satisfied [18].

Combining Eqs. (32), (33) and (36) we find

$$\delta_H^2 \approx \frac{\sqrt{3}}{75\,\pi^2}\,\frac{\exp[-2\widetilde{\Im}(\phi)]}{r^{1/2}\,\widetilde{\varepsilon}}\,\frac{T_r}{H}, \tag{37}$$

or equivalently by using Eqs.(9) and (10) we obtained

$$\delta_H^2 \approx \frac{\sqrt{3}}{75\,\pi^2}\exp[-2\widetilde{\Im}(\phi)]\left[\left(\frac{1}{\widetilde{\varepsilon}}\right)^3\,\frac{1}{2\,r^2\,\sigma\,\kappa^2\,V}\right]^{1/4}, \tag{38}$$

where the dimensionless slow-roll parameter in the hight dissipation period becomes

$$\widetilde{\varepsilon} \approx \frac{1}{6\kappa\, r}\left[\frac{V_{,\phi}}{V}\right]^2\frac{1}{V}\;. \tag{39}$$

The scalar spectral index n_s is given by

$$n_s - 1 = \frac{d\ln\,\delta_H^2}{d\ln k}, \tag{40}$$

where the interval in wavenumber is related to the number of e-folds by the relation $d\ln k(\phi) = -dN(\phi)$, and using Eqs.(38) and (40), it becomes

$$n_s \approx 1 - \left[\frac{3\widetilde{\eta}}{2} + \widetilde{\varepsilon}\left(\frac{2V}{V_{,\phi}}\left[2\widetilde{\Im}_{,\phi} - \frac{r_{,\phi}}{4r}\right] - \frac{5}{2}\right)\right], \tag{41}$$

where the second slow-roll parameter η (for $r \gg 1$) is given by

$$\widetilde{\eta} \approx \frac{1}{3\kappa\, r\, V}\left[\frac{V_{,\phi\phi}}{V} - \frac{1}{2}\left(\frac{V_{,\phi}}{V}\right)^2\right]. \tag{42}$$

One of feature of the three-year data set from WMAP is that it hints at a significant running in the scalar spectral index $dn_s/d\ln k = \alpha_s$ [4]. Dissipative effects themselves can produce a rich variety of spectral ranging between red and blue [7, 15]. From Eq.(41) we obtain that the running of the scalar spectral index for our model becomes

$$\alpha_s = \frac{dn_s}{d\ln k} \approx \frac{2\,V\,\widetilde{\varepsilon}}{V_{,\,\phi}}\left[\frac{3\widetilde{\eta}_{,\,\phi}}{2} + \frac{\widetilde{\varepsilon}_{,\,\phi}}{\widetilde{\varepsilon}}\left(n_s - 1 + \frac{3\widetilde{\eta}}{2}\right) + 2\widetilde{\varepsilon}\left(\left(\frac{V}{V_{,\,\phi}}\right)_{,\,\phi}\left[2\widetilde{\Im}_{,\,\phi} - \frac{(\ln r)_{,\,\phi}}{4}\right]\right.\right.$$

$$\left.\left. + \left(\frac{V}{V_{,\,\phi}}\right)\left[2\widetilde{\Im}_{,\,\phi\,\phi} - \frac{(\ln r)_{,\,\phi\,\phi}}{4}\right]\right)\right]. \qquad (43)$$

In models with only scalar fluctuations, the marginalized value for the derivative of the spectral index is approximated as $dn_s/d\ln k = \alpha_s \sim -0.05$ for WMAP-3 only [4].

In the next section we will study the specific cases in which the dissipation parameter, Γ, is either $\Gamma = const.$ or $\Gamma = \Gamma(\phi)$.

III-2. Tensor Perturbations

As it is mentioned in Ref.[19], the generation of tensor perturbation during inflation would produce stimulated emission in the thermal background of gravitational wave. This process changes the power spectrum of the tensor modes through an extra temperature dependence factor $\coth(k/2T)$. The spectrum in this case is given by

$$A_g^2 = \frac{16\pi}{m_p^2}\left(\frac{H}{2\pi}\right)^2 \coth\left[\frac{k}{2T}\right] \simeq \frac{32\,V}{3\,m_p^4}\coth\left[\frac{k}{2T}\right], \qquad (44)$$

where the spectral index n_g results as

$$n_g = \frac{d}{d\,\ln k}\,\ln\left[\frac{A_g^2}{\coth[k/2T]}\right] = -2\,\varepsilon. \qquad (45)$$

Here, we have used that $A_g^2 \propto k^{n_g}\,\coth[k/2T]$ and Eq.(11) [19].

From expressions (37), and (44), we may write the tensor-scalar ration as

$$R(k_0) = \left(\frac{A_g^2}{P_{\mathcal{R}}}\right)\Bigg|_{k=k_0} = \frac{240\sqrt{3}}{25m_p^2}\left[\frac{(1+r)\,\varepsilon\,H^3}{r^{1/2}\,T_r}\,\exp[2\,\Im(\phi)]\,\coth\left(\frac{k}{2T}\right)\right]\Bigg|_{k=k_0},$$

where we have used that $\delta_H \equiv 2\,P_{\mathcal{R}}^{1/2}/5$. For $r \gg 1$, this relation reduces to

$$R(k_0) \approx \frac{240\sqrt{3}}{25m_p^2}\left[\frac{r^{1/2}\,\widetilde{\varepsilon}\,H^3}{T_r}\,\exp[2\,\widetilde{\Im}(\phi)]\,\coth\left(\frac{k}{2T}\right)\right]\Bigg|_{k=k_0}, \qquad (46)$$

where k_0 is referred to as the pivot point.

From the combination of WMAP three-year data[4] with the SDSS large scale structure surveys [20], there is found an upper bound $R(k_0{=}0.002\ \mathrm{Mpc}^{-1}) < 0.28(95\%CL)$, where $k_0{=}0.002\ \mathrm{Mpc}^{-1}$ corresponds to $l = \tau_0 k_0 \simeq 30$, with the distance to the decoupling surface $\tau_0 =$ 14,400 Mpc. SDSS measures galaxy distributions at red-shifts $a \sim 0.1$ and probes k in the range 0.016 h Mpc^{-1}< k <0.011 h Mpc^{-1}. The recent WMAP three-year results give the values for the scalar curvature spectrum $P_{\mathcal{R}}(k_0) \equiv 25\delta_H^2(k_0)/4 \simeq 2.3 \times 10^{-9}$ and the scalar-tensor ration $R(k_0) = 0.095$. These values allow us to find constrains on the parameters of our model.

IV. EXPONENTIAL POTENTIAL IN THE HIGH DISSIPATION APPROACH.

The tachyonic effective potential $V(\phi)$ is one that satisfies $V(\phi) \longrightarrow 0$ as $\phi \longrightarrow \infty$. It has been argued that the qualitative tachyonic potential of string theory can be describe via an exponential potential of the form [11]

$$V(\phi) = V_0 e^{-\alpha\phi}, \tag{47}$$

where α and V_0 are free parameters. In the following we will take $\alpha > 0$ (with unit m_p). The parameter α is related to the tachyon mass [21]. An estimation of these parameters is give for the cool inflationary case in [12], where $V_0 \sim 10^{-10} m_p^4$ and $\alpha \sim 10^{-6} m_p$.

In the following we will restrict ourselves to the high dissipation regimen in which $r \gg 1$.

IV-1. $\Gamma = \Gamma_0 = const.$ case

With $\Gamma = \Gamma_0 = const.$ and using the exponential potential given by Eq.(47), we find that the slow roll parameters becomes

$$\widetilde{\varepsilon} = \widetilde{\eta} = \frac{1}{6\kappa\, r} \frac{\alpha^2}{V_0 e^{-\alpha\phi}}. \tag{48}$$

The Hubble parameter is given by

$$H(\phi) = \sqrt{\kappa V_0}\, e^{-\alpha\phi/2}, \tag{49}$$

while the rate r becomes

$$r = \frac{m_p \Gamma_0}{\sqrt{24\pi}} \frac{1}{V_0^{3/2}}\, e^{3\alpha\phi/2} \gg 1. \tag{50}$$

The evolution of the $\dot{\phi}$ during this scenario is governed by the expression

$$\dot{\phi} = -\frac{V_{,\phi}}{3rHV}, \tag{51}$$

and using Eqs.(49)-(51), we find that the evolution of the tachyonic field as a function of time is given by

$$\phi(t) = \frac{1}{\alpha}\,\ln\left[e^{\alpha\phi_i} + \frac{\alpha^2 V_0}{\Gamma_0}\,t\right], \tag{52}$$

where $\phi(t = t_i = 0) = \phi_i$. Substituting this last equation into Eq.(49), we obtained for the scale factor

$$\frac{a(t)}{a_i} = \exp\left[\frac{2\Gamma_0}{\alpha^2}\sqrt{\kappa/V_0}\left[\sqrt{e^{\alpha\phi_i} + \frac{\alpha^2 V_0}{\Gamma_0}\,t} - e^{\alpha\phi_i/2}\right]\right]. \tag{53}$$

It is not hard to see that $\ddot{a}(t = t_f) = 0$ (or equivalently $\epsilon_f \simeq 1$), where we get that

$$t_f = \frac{\alpha^2}{4\kappa\,\Gamma_0} - \frac{\Gamma_0\,e^{\alpha\phi_i}}{\alpha^2\,V_0}. \tag{54}$$

Thus, t_f represents the time at which inflation ends. Substituting Eq.(79) into Eq.(52),we find that the tachyonic field at the end of inflation becomes

$$\phi_f = -\frac{1}{\alpha}\,\ln\left[\frac{4\kappa\,\Gamma_0^2}{\alpha^4 V_0}\right]. \tag{55}$$

The energy density of the radiation field becomes

$$\rho_\gamma = \frac{\sqrt{6\pi}}{6\kappa}\left[\frac{\alpha^2}{\Gamma_0}\right]\,V_0^{3/2}\,e^{-3\alpha\phi/2}, \tag{56}$$

and, in terms of ρ_ϕ, it is given by

$$\rho_\gamma = \frac{\sqrt{6\pi}}{6\kappa}\left[\frac{\alpha^2}{\Gamma_0}\right]\,\rho_\phi^{3/2}. \tag{57}$$

Using Eq. (16), the total number of e-folds at the end of warm inflation results as

$$N_{total} = -3\kappa\int_{\phi_i}^{\phi_f}\frac{V^2}{V_{,\tilde{\phi}}}\,r\,d\tilde{\phi} = \frac{2\Gamma_0}{\alpha^2}\left[\frac{\kappa}{V_0}\right]^{1/2}[\exp(\alpha\phi_f/2 - \exp(\alpha\phi_i/2)], \tag{58}$$

where the initial tachyonic field satisfies $\phi_i < \phi_f$, since $V_i > V_f$.

Rewriting the total number of e-folds in terms of V_f and V_i, and using Eq.(55), we find

$$N_{total} = \left[1 - \left(\frac{V_f}{V_i}\right)^{1/2}\right]. \tag{59}$$

Note that $V_i > V_f$ and thus we obtain $N_{total} \lesssim 1$ e-fold. This problem it is not surprising since this lies in the slow roll approximation which by virtue of Eq.(48) is valid when $\phi \ll \ln r$, i.e., when ϕ is not too large for the case in which $r \gg 1$. This implies, from Eq.(52), that this solution is valid only at early times and therefore it is not asymptotic. Therefore, the problem of small e-folds is due to the slow roll approximation. In section V-1, we will again consider the case $\Gamma = const.$, but there a power law scale factor is assumed.

IV-2. Γ as a function of ϕ.

In this case we consider Γ as a function of ϕ. We take this to be of the form, $\Gamma = f(\phi) = c^2\, V(\phi) = c^2\, V_0\, e^{-\alpha\phi}$, where $c^2 > 0$ (with unit of m_p). The slow roll parameters are equal, i.e.

$$\widetilde{\varepsilon} = \widetilde{\eta} = \frac{\alpha^2\, e^{\alpha\phi/2}}{2\, c^2\, (\kappa\, V_0)^{1/2}}, \tag{60}$$

and the dissipation parameter r is given by

$$r = \frac{c^2\, e^{\alpha\phi/2}}{3(\kappa\, V_0)^{1/2}}. \tag{61}$$

In this case, we obtain that the evolution of the tachyon field is given by

$$\phi(t) = \phi_i + \frac{\alpha}{c^2}\, t, \tag{62}$$

and the scale factor becomes

$$\frac{a(t)}{a_i} = \exp\left[\frac{2c^2}{\alpha^2}\sqrt{\kappa V_0}\, e^{-\alpha\phi_i/2}[1 - e^{-\alpha^2\, t/2c^2}]\right]. \tag{63}$$

The value of the potential at the end of inflation is given by

$$V_f = V_0 e^{-\alpha\phi_f} = \frac{\alpha^4}{4\,\kappa\, c^4}, \tag{64}$$

and the total number of e-folds is related to V_f and V_i through

$$V_i = (2\, N_{total} - 1)^2\, V_f. \tag{65}$$

Note that, unlike in the previous case, the N_{total} parameter could assume appropriate values (60 or so) for solving the standard cosmological puzzles. To do that,we need the following inequality to be satisfied: $V_i > 10^4\, V_f$.

At the beginning of inflation the dissipation parameter becomes

$$r(\phi) = r_i = \frac{2}{3}\frac{c^4}{(2\,N_{total}-1)\,\alpha^2} \gg 1, \tag{66}$$

resulting in the requirement that $c^2 \gg \alpha\,(N_{total}-1)^{1/2}$.

On the other hand, from Eq. (30) we obtain that

$$\delta\phi \simeq C\frac{\dot{\phi}}{H}\left[1+\frac{c^2}{3H}\right]^{\frac{5}{4}+\frac{21\alpha^2}{16c^4}} \exp\left[\frac{c^2(6H^2(4c^4-21\alpha^2)+Hc^2(8c^4-63\alpha^2)-6c^4\alpha^2)}{32H(c^2+3H)^2}\right]. \tag{67}$$

From Eq.(23), the metric perturbation Φ becomes given by

$$\Phi \simeq -C_1\frac{\dot{H}}{H^2}\left[1+\frac{c^2}{3H}\right]^{\frac{5}{4}+\frac{21\alpha^2}{16c^4}} \left[1+\frac{c^2}{4H}-\frac{\alpha^2\,c^2}{48H^2(c^2+3H)}\right]$$

$$\times\;\exp\left[\frac{c^2(6H^2(4c^4-21\alpha^2)+Hc^2(8c^4-63\alpha^2)-6c^4\alpha^2)}{32H(c^2+3H)^2}\right], \tag{68}$$

where the zeroth-order term is related the adiabatic mode and the remaining terms describe the inclusion of dissipation in the fluctuation of the scalar field (the entropy mode)[16].

From Eq.(40) we obtained that the scalar spectral index of the adiabatic perturbation becomes

$$n_s \approx 1-\left[\frac{3\,\widetilde{\eta}}{2}-\frac{3}{4}\,\widetilde{\varepsilon}(1+\widetilde{\varepsilon})\right], \tag{69}$$

which becomes r dependent for the case of high dissipation.

From Eq.(37), when $r \gg 1$ or equivalently $c^2 \gg 3H$, we obtain that the scalar power becomes

$$P_{\mathcal{R}}(k_0) \approx \frac{\kappa^{19/4}}{2\,\pi^2}\left[\frac{c}{\alpha^2}\,V(\phi_0)^{5/2}\,T_r\right]\exp\left[\frac{3\alpha^2}{8c^2\sqrt{\kappa\,V(\phi_0)}}\right], \tag{70}$$

and from Eq.(46) the tensor-scalar ration is given by

$$R(k_0) \approx 3\,\pi^2\left(\frac{\alpha^2}{c\,\kappa^{11/4}}\right)\left[\frac{1}{V(\phi_0)^{3/2}\,T_r}\right]\exp\left[\frac{-3\alpha^2}{8c^2\sqrt{\kappa\,V(\phi_0)}}\right]\coth\left[\frac{k_0}{2T}\right], \tag{71}$$

where $V(\phi_0) = V_0\exp[-\alpha\phi_0]$ and ϕ_0 represents the value of the tachyon field when the scale $k_0 = 0.002\text{Mpc}^{-1}$ was leaving the horizon.

Using the WMAP three-year data where $P_{\mathcal{R}}(k_0) \simeq 2.3\times10^{-9}$, $R(k_0) = 0.095$ and choosing the parameters $c^2 = 10^{10}m_p$, $T \simeq T_r \simeq 0.24\times10^{16}$ GeV and $k_0 = 0.002\text{Mpc}^{-1}$, we obtained from Eqs.(70) and (71) that $V(\phi_0) \simeq 2.43\times10^{-15}m_p^4$ and $\alpha \simeq 1.76\times10^{-12}\,m_p$. We noted that the dissipation coefficient when the scale k_0 was leaving the horizon is the order of $\Gamma(k_0) \sim 10^{-5} m_p^5$. From Eq.(43) we found that it is necessary to increase the value of α by several orders of magnitude in order to have a running spectral index α_s close to value given by WMAP observations.

V. POWER LAW IN THE HIGH DISSIPATION APPROACH.

An interesting approach is to choose a scale factor of power law type, i.e. $a \sim t^p$, in which $p \geq 2$. This allows us to get the form of the scalar potential $V(\phi)$ for different values of Γ in the high dissipation approach, i.e. $\tilde{r} = \frac{\Gamma}{3H\rho_\phi} \gg 1$. Note that if $\dot{\phi}^2 \ll 1$, $\rho_\phi \sim V$ we get $\tilde{r} \longrightarrow r$.

In the following we will consider some special cases for the parameter Γ.

V-1. $\Gamma = \Gamma_0 = const.$ case

By using Eqs.(4) and (5), we obtained that

$$\dot{\rho}_\phi + \dot{\rho}_\gamma + 4\ H\ \rho_\gamma = 0, \tag{72}$$

where we have used for the energy density and pressure associated with the tachyon field the relation $p_\phi = -\rho_\phi\,(1 - \dot{\phi}^2)$. From Eqs.(3) and (72) and choosing a scale factor of power law type, we find that

$$\rho_\gamma = \frac{p}{2\kappa\, t^2}\ , \quad \rho_\phi = \frac{p\ (2p-1)}{2\kappa\, t^2} = \rho_\gamma(2p-1). \tag{73}$$

From Eq.(4)(or equivalently from Eq.(5)) the temporal evolution of the tachyonic field becomes given by

$$(\phi - \phi_1)^2 = \frac{4(2p-1)}{\kappa\Gamma_0}\,\frac{p}{t}, \tag{74}$$

where ϕ_1 is a constant of integration. Substituting Eq.(73) into Eq.(1) and using Eq.(74), we obtained that the potential as a function of the tachyonic field ϕ becomes

$$V(\phi) = \beta_1\,(\phi - \phi_1)^4\,\sqrt{1 - \beta_2\,(\phi - \phi_1)^6}\,, \tag{75}$$

where β_1 and β_2 are given by

$$\beta_1 = \frac{\kappa\Gamma_0^2}{32p\,(2p-1)}\ ,\ \mathit{and}\quad \beta_2 = \frac{\kappa^2\Gamma_0^2}{64\,p^2\,(2p-1)^2},$$

respectively.

We note that Eq. (72) is valid for

$$\tilde{r} = \frac{\Gamma}{3H\rho_\phi} \gg 1 \implies \frac{128\, p\,(2p-1)^2}{3\kappa^2\,\Gamma_0^2} \gg (\phi-\phi_1)^6. \tag{76}$$

The number of e-folds becomes

$$N = \int_t^{t_f} H(t')\,dt' = 2\,p\,\ln\left[\frac{\phi-\phi_1}{\phi_f-\phi_1}\right], \tag{77}$$

where the time at the end of inflation is calculated when $\ddot{a}=0$, or equivalently, by using the relation

$$[\rho_\phi + \rho_\gamma = -3(p_\phi + p_\gamma)]|_{\phi(t=t_f)}\ , \tag{78}$$

which gives $\dot{\phi}_f^2 = 1-(2p+1)/(3(2p-1))$.

From Eq.(74) we find that the time at the end of inflation becomes given by

$$t_f = \left[\frac{3\,p\,(2p-1)^2}{4\kappa\Gamma_0(p-1)}\right]^{1/3}, \tag{79}$$

and from Eqs.(77) and (79), we have that

$$(\phi-\phi_1) = \left[\frac{4^4\,p^2(2p-1)(p-1)}{3\kappa^2\Gamma_0^2}\right]^{1/6}\ \exp\left[\frac{N}{2p}\right]. \tag{80}$$

On the other hand, the spectrum of scalar perturbations becomes

$$P_{\mathcal{R}}(k_0) \approx \frac{2}{\pi^2}\left[\frac{V(\phi_0)^{23}}{\beta_1^{37/2}}\right]^{1/4}\left(\frac{T_r}{(\phi_0-\phi_1)^{13}\,[4-7\beta_2(\phi_0-\phi_1)^6]^2}\right), \tag{81}$$

and the tensor-scalar ration is given by

$$R(k_0) \approx \frac{512\pi}{m_p^2}\left[\frac{\beta_1^{53/2}}{V(\phi_0)^{23}}\right]^{1/4}\left(\frac{(\phi_0-\phi_1)^{17}\,[4-7\beta_2(\phi_0-\phi_1)^6]^2}{\Gamma_0^2\,T_r}\right)\ \coth\left[\frac{k_0}{2T}\right]. \tag{82}$$

Using the WMAP three-year data, in which $P_{\mathcal{R}}(k_0) \simeq 2.3\times10^{-9}$, $R(k_0) = 0.095$ and choosing the parameters $p=2$, $T \simeq T_r \simeq 0.24\times10^{16}$ GeV and $k_0 = 0.002\text{Mpc}^{-1}$, we obtain from Eqs.(81) and (82) that $(\phi_0-\phi_1) \approx 2\times10^3 m_p$ and $\Gamma_0 \approx 10^{-11}m_p^5$. Here the scalar potential when the scale is $k_0 = 0.002\text{Mpc}^{-1}$ is leaving the horizon becomes $V(\phi_0) \sim 3\times10^{-10}m_p^4$. Note that this value of $V(\phi_0)$ becomes similar to that found for cool inflation where a chaotic potential is used [19]. Another case of interest is that of $p=4$, where it is found that $(\phi_0-\phi_1) \approx 10^7 m_p$, $\Gamma_0 \approx 10^{-22}m_p^5$ and $V(\phi_0) \sim 10^{-17}m_p^4$. In this case, from Eq.(43) we found that if we decrease the value of $(\phi-\phi_1)$ by a few orders of magnitude, then the running spectral index α_s takes a value which agrees with the value specified by the WMAP observations.

V-2. Γ as a function of t

In this case, for simplicity we considered that the dissipation coefficient is $\Gamma = \gamma/t$, where γ is a positive-definite constant (with the unit of m_p^4). By using Eqs.(4) and (73), we obtain that the tachyonic field becomes

$$(\phi - \tilde{\phi}_1) = \beta_3 \, \ln[t], \tag{83}$$

where $\tilde{\phi}_1$ is a constant of integration and $\beta_3 = \sqrt{(2p-1)\, p/\kappa\gamma}$. We also note that Eq. (72) is valid for

$$\tilde{r} = \frac{\Gamma}{3H\rho_\phi} \gg 1 \implies \frac{2\kappa\gamma}{3\,p^2\,(2p-1)} \gg \exp(-2(\phi - \phi_1)/\beta_3). \tag{84}$$

The potential is now given by

$$V(\phi) = \left(\frac{\gamma\,\beta_3^2}{2}\right) \exp\left[\frac{-2(\phi - \tilde{\phi}_1)}{\beta_3}\right] \sqrt{1 - \beta_3^2 \, \exp\left[\frac{-2(\phi - \tilde{\phi}_1)}{\beta_3}\right]}. \tag{85}$$

From Eqs.(78) and (83), we obtain that the time at the end of inflation is given by

$$t_f = \frac{(2p-1)}{2} \sqrt{\frac{3\,p}{\kappa\,\gamma\,(p-1)}}\,. \tag{86}$$

The tachyonic field in terms of the N e-folds parameter becomes

$$(\phi - \tilde{\phi}_1) = \beta_3 \left[\ln\left(\frac{(2p-1)}{2} \sqrt{\frac{3\,p}{\kappa\,\gamma\,(p-1)}}\right) - \frac{N}{p}\right]. \tag{87}$$

Following a scheme similar to those of the previous cases, in order to calculate the spectrum of scalar perturbations and the tensor-scalar ratio, we obtained that $(\phi_0 - \tilde{\phi}_1) \approx 69.6 m_p$ and $\gamma \approx 0.02\, m_p^4$, and the potential when the scale $k_0 = 0.002\text{Mpc}^{-1}$ is leaving the horizon becomes $V(\phi_0) \sim 0.2 \times 10^{-14}\, m_p^4$. We have chosen the values $p = 2$ and $T \simeq T_r \simeq 0.24 \times 10^{16}$ GeV. For the value of the parameters given above, and from Eq.(43), we find that the running spectral index α_s lies in the range established by WMAP observations.

VI. CONCLUSIONS

In this paper we have investigated the tachyonic warm inflationary scenario. In the slow roll approximation we have found a general relationship between the radiation and tachyon energy densities. This has led us to a general criterion for tachyon warm inflation to occur; see Eq.(14).

In relation to the corresponding perturbations, the contributions of the adiabatic and entropy modes were obtained explicitly. Here, it is shown that the dissipation parameter plays a crucial role in producing the entropy mode. Additionally, we have found a general relation for the density perturbation expressed by Eq.(32). The tensor perturbation are generated via stimulated emission into the existing thermal background (see Eq.(44)), and the tensor-scalar ratio is modified by a temperature dependent factor.

Our first specific models are described using an exponential scalar potential and for two different dependences of the dissipation coefficient, Γ. In the first case, we took $\Gamma = \Gamma_0 = Cte.$ and we have found that the number of e-folds in this case is insufficient. For the case in which the dissipation coefficient Γ is taken to be a function of the tachyon field, i.e. $\Gamma = f(\phi) = c^2 \ V(\phi)$, it has been possible to describe appropriately tachyon warm inflationary universe models. Also, in place of choosing a specific form of the scalar tachyon potential a priory, we have assumed a power law for the scale factor. Here, we have described inflationary models for $\Gamma = \Gamma_0 = const$ and for Γ variable. In both cases, we have obtained explicit expressions for the corresponding scalar potential. By using the WMAP three-year data, we have found some constraints for the parameters appearing in our model. From the normalization of the WMAP data, the potential becomes of the order of $V(\phi_0) \sim 10^{-15} m_p^4$ when it leaves the horizon, at the scale of $k_0 = 0.002\mathrm{Mpc}^{-1}$.

Dissipative effects plays a crucial role in producing the entropy mode; they can themselves produce a rich variety of spectral ranging between red and blue. The possibility of a spectrum which runs from blue to red is particularly interesting because it is not commonly seen in inflationary models, which typically predict red spectral. Models of inflation with dissipative effects and models with interacting fields have much more freedom to yield spectral. In particular, we anticipate that the Planck mission will significantly enhance our understanding of α_s by providing hight quality measurements of the fundamental power spectrum over a large wavelength range than WMAP. Summarizing, we have been successful in describing tachyon warm inflationary models for describing the early epoch of the universe in the high dissipation regimen.

Acknowledgments

R. H. was supported by the “Programa Bicentenario de Ciencia y Tecnología” through the Grant “Inserción de Investigadores Postdoctorales en la Academia” N^0 PSD/06. S. d. C. was supported by COMISION NACIONAL DE CIENCIAS Y TECNOLOGIA through FONDECYT grants N^0 1030469, $N^0$1040624 and $N^0$1051086. Also, from UCV-DGIP N^0 123.764. C. C. was supported by MINISTERIO DE EDUCACION through Postdoctoral MECESUP Grants FSM 0204.

[1] Guth A., The inflationary universe: A possible solution to the horizon and flatness problems, 1981 Phys. Rev. D **23**, 347.

[2] Albrecht A. and Steinhardt P. J., Cosmology for grand unified theories with radiatively induced symmetry breaking, 1982 Phys. Rev. Lett. **48**, 1220; A complete description of inflationary scenarios can be found in the book by Linde A., *Particle physics and inflationary cosmology* (Gordon and Breach, New York, 1990).

[3] Spergel D. N., First year Wilkinson microwave anisotropy probe (WMAP) observations: Determination of cosmological parameters, 2003 Astrophys. J. Suppl. **148**, 175; Peiris H.V., First year wilkinson microwave anisotropy probe (WMAP) observations: Implications for inflation, 2003 Astrophys. J. Suppl. **148**, 213.

[4] Page L. et al., Wilkinson microwave anisotropy probe (WMAP) three year results: implications for cosmology. astro-ph/0603449.

[5] Linde A., A new inflationary universe scenario: a possible solution of the horizon, flatness, homogeneity, isotropy and primordial monopole problems, 1982 Phys.Lett. B **108**, 389; Albrechtand A., Steinhart P.J., 1982 Cosmology for grand unified theories with radiatively induced symmetry breaking 1982 Phys.Rev.Lett. **48**, 1220; Hawking S.W., Moss I.G., Supercooled phase transitions in the very early universe, 1982 Phys.Lett.B **110**, 35.

[6] Berera A. , Warm inflation, 1995 Phys. Rev. Lett. **75**, 3218; Berera A., Interpolating the stage of exponential expansion in the early universe: A possible alternative with no reheating, 1997 Phys. Rev. D **55**, 3346.

[7] Hall L.M.H., Moss I.G. and Berera A. Scalar perturbation spectra from warm infaltion, 2004

Phys.Rev.D **69**, 083525; Moss I.G. Primordial inflation with spontaneous symmetry breaking, 1985 Phys.Lett.B **154**, 120; Berera A., Fang L.Z., Thermally induced density perturbatons in the inflatiionary era, 1995 Phys.Rev.Lett. **74** 1912; Berera A., Warm inflation at arbitrary adiabaticity: A model, an existence proof for inflationery dynamics in quantum field theory, 2000 Nucl.Phys B **585**, 666.

[8] Berera A., Thermal properties of an inflationary universe, 1996 Phys. Rev.D **54**, 2519.

[9] Taylor A.N. and Berera A., Perturbation spectra in the warm inflationary scenario, 1997 Phys. Rev. D **55**, 3346.

[10] Sen A., Rolling tachyon, 2002 JHEP **0204**, 048 .

[11] Sen A., Field theory of tachyon matter, 2002 Mod. Phys. Lett. A **17**, 1797 .

[12] Sami M., Chingangbam P. and Qureshi T, Aspects of tachyonic inflation with exponential potential, 2002 Phys. Rev. D **66**, 043530.

[13] Gibbons G. W., Cosmological evolution of the rolling tachyon, 2002 Phys. Lett. B **537**, 1.

[14] Bardeen J., Gauge Invariant Cosmological Perturbations, 1980 Phys. Rev. D **22**, 1882.

[15] Starobinsky A. and Yokoyama J., Density fluctuations in Brans-Dicke inflation, Published in the Proceedings of the Fourth Workshop on General Relativity and Gravitation. Edited by K. Nakao, et al. Kyoto, Kyoto University, 1995. pp. 381, gr-qc/9502002; Starobinsky A. and Tsujikawa S., Cosmological perturbations from multifield inflation in generalized Einstein theories, 2001 Nucl.Phys.B **610**, 383; Oliveira H. and Joras S., On perturbations in warm inflation, 2001 Phys. Rev. D **64**, 063513.

[16] Oliveira H., Density perturbations in warm inflatio and COBE normalization, 2002 Phys. Lett. B **526**, 1.

[17] Liddle A. and Lyth D., Cosmological inflation and large-scale structure, 2000, Cambridge University; Linsey J., Liddle A., Kolb E. and Copeland E., 1997 Rev. Mod. Phys **69**, 373; Bassett B., Tsujikawa S. and Wands D., Inflation dynamics and reheating, 2005 Rev. Mod. Phys. **78**, 537.

[18] Taylor A. and Berera A., Perturbations spectral in the warm inflationary scenario, 2000 Phys. Rev. D **62**, 083517.

[19] Bhattacharya K., Mohanty S. and Nautiyal A., Enhanced polarization of CMB from thermal gravitational waves, astro-ph/0607049.

[20] Tegmark M. et al., Cosmological parameters from SDSS and WMAP, 2004 Phys. Rev. D **69**,

103501.

[21] Fairbairn M. and Tytgat M., Inflation from a tachyon fluid?, 2002 Phys. Lett. B **546**, 1.